

Detailed Description of the Invention

As discussed supra, while many groups have previously reported the production of anti-CD23 antibodies and the use thereof as antagonists and agonists for modulating IgE production, the exact mechanism by which such antibodies modulate IgE expression in systems where IL-4 induces IgE production remains unclear. Thus, it would be beneficial if the means by which such antibodies modulate IgE expression were elucidated, or at least better explained, as such information would be potentially useful in designing therapeutics for treatment of diseases wherein modulation of IgE production is therapeutically desirable. In particular, it would be beneficial if improved antibodies specific to CD23 were obtained having improved capacity to inhibit induced IgE production, as enhanced IgE levels are believed to be involved in numerous disease processes, e.g., allergic conditions, inflammatory conditions and autoimmune diseases. Such diseases include by way of example, atopic dermatitis, eczema, allergic rhinitis and conjunctivitis, Job's syndrome, and asthma.

Toward that end, the present inventors have surprisingly discovered that anti-human CD23 monoclonal antibodies which contain human gamma-1 constant domains inhibit IgE production in systems where IgE production is induced by IL-4 significantly better than CD23 monoclonal antibodies of other effector types, e.g., those comprising human

gamma-4 constant domains or CD23 monoclonal antibodies or antibody fragments lacking effector functions altogether.

While Fc mediated effector functions are sometimes significant to the therapeutic activity of antibodies, this discovery was surprising in the case of anti-CD23 antibodies because the role of effector function in the IgE inhibitory activity of anti-CD23 antibodies had not been previously reported. In fact, previous evidence had suggested that antibody effector function was not significant to the ability of anti-CD23 antibodies to inhibit induced IgE production. For example, Flores-Romo et al., *Science*, 261:1038-1041 (1993) had reported that Fabs prepared from a polyclonal anti-CD23 antibody inhibited an *in vivo* induced IgE antigen-specific response.

This discovery was made after the present inventors isolated various primate antibodies specific to CD23 having anti-IgE inhibiting activity and compared these antibodies to PRIMATIZED® versions with respect to their ability to inhibit IL-4 induced IgE production *in vitro* and *in vivo*.

More specifically, and as described in greater detail *infra*, five primate monoclonal antibodies which specifically bound both cellular and soluble CD23 were isolated from an Old World monkey (macaque) according to the methodology which is disclosed in <sup>U.S. Patent No. 5,058,570</sup> ~~commonly assigned Application Serial No. 08/379,072 (now allowed)~~, which applica-

tion is incorporated by reference in its entirety herein. This application described in detail a means for producing monoclonal antibodies to desired antigens, desirable human antigens, in Old World monkeys and their advantages in relation to antibodies of other species as therapeutics, for example reduced or potentially lack of immunogenicity in humans because of the phylogenetic closeness of humans and Old World monkeys. In fact, because of the phylogenetic closeness of these species, it is difficult to distinguish Old World monkey immunoglobulins from human immunoglobulins by sequence comparison.

Four of these five primate monoclonal anti-human CD23 antibodies were demonstrated to be capable of inhibiting IL-4 induced IgE production in an *in vitro* B cell assay described in detail *infra* and the most potent was also shown to inhibit IL-4 induced IgE in a SCID mouse animal model (also described in detail *infra*). Based on this IgE inhibitory activity, and expected low immunogenicity in humans, such antibodies are potentially suitable as therapeutics for treating diseases wherein inhibition of IgE production is therapeutically desirable.

However, in order to further reduce immunogenicity, it was elected to PRIMATIZE® two primate monoclonal antibodies (a type of chimerization of antibodies) according to the methodology which is also described in U.S. Serial No. 08/379,072 (now allowed), incorporated by reference

herein. PRIMATIZATION® essentially refers to the production of recombinant antibodies developed by IDEC Pharmaceuticals Corporation which comprise primate variable regions and human constant regions. Primatization of the two primate anti-human CD23 monoclonal (5E8 and 6G5) antibodies having potent IgE inhibiting activity was effected in order to eliminate any potential immunogenicity attributable to the primate constant domains in humans.

Because of the inventors' initial expectation from published literature that Fc effector function was not necessary for induced IgE inhibition, human gamma 4 versions of these particular antibodies were initially produced. However, quite surprisingly, it was found that the gamma-4 versions produced from both of these primate monoclonal antibodies were ineffective, i.e., they required significantly higher concentrations of PRIMATIZED® gamma 4 antibody than the primate antibody to inhibit IL-4 induced IgE production in in vitro assays.

Moreover, even more surprising was the discovery that when the same two primate antibodies were then converted to human gamma-1 versions (by substitution of the primate constant domains with human gamma-1 constant domains), that these gamma-1 antibodies very effectively inhibited induced IgE production in vitro. Thus, our results suggested that Fc effector function is apparently significant

to the ability of anti-human CD23 antibodies to inhibit induced IgE production. This hypothesis was confirmed when a third primate anti-human CD23 monoclonal, i.e., the 2C8 antibody, which was shown by us to inhibit IgE production in vitro, was converted to a F(ab')<sub>2</sub>, which was found to be substantially incapable of inhibiting induced IgE production in vitro. In fact, this F(ab')<sub>2</sub> was found to antagonize the suppressive effects on induced IgE blocking activity of the primate anti-human CD23 monoclonal antibody 2C8.

In addition, it was found that removing a glycosylation site in the heavy chain variable region of one of the antibodies (5E8) had no effect on binding of the antibody to CD23 (as evidenced by obtained K<sub>d</sub> values), or on induced IgE inhibition. Thus, the differences in IgE inhibition were shown to apparently not involve glycosylation differences.

The PRIMATIZED® gamma 1 version of primate 6G5 was found to inhibit induced IgE expression in SCID mice while the same concentration of either the primate 6G5 or the PRIMATIZED® p6G5F4p did not inhibit induced IgE expression. Therefore, an antibody containing human gamma-1 constant domains was found to be even more effective in an in vivo animal model than the primate monoclonal antibody.

Accordingly, based on these results, it has been surprisingly discovered that an active Fc region, in particu-

lar that of human gamma 1, is significantly involved in the mechanism of IL-4 induced IgE inhibition by anti-human CD23 monoclonal antibodies. This discovery is quite unexpected especially based on earlier reports that Fabs derived from polyclonal anti-CD23 antibodies were capable of inhibiting induced IgE production, and also based on the various theories as to how CD23 affects induced IgE expression.

Accordingly, the present invention relates to anti-human CD23 antibodies containing human gamma-1 constant domains and their use as therapeutics based on their ability to effectively inhibit IgE expression.

The skilled artisan can prepare anti-human CD23 antibodies containing human gamma-1 constant domains by methods which are well known in the art for the manufacture of chimeric antibodies. Essentially, such methods comprise producing anti-human CD23 antibodies in a desired host or *in vitro*, cloning a hybridoma or cell line which produces an anti-human CD23 monoclonal antibody exhibiting desirable characteristics, e.g., adequate CD23 binding affinity, cloning the nucleic acid sequences which encode such antibody from said hybridoma or cell line, e.g. by polymerase chain reaction using suitable primers, isolating the variable domains contained therein, recombining such variable domains with human gamma-1 constant domains and the appropriate human light chain constant domain, and

- 20 -

expressing the resultant nucleic acid sequence encoding a chimeric anti-human CD23 gamma-1 immunoglobulin in a suitable expression system. Preferably, the anti-human CD23 antibodies of the invention will have apparent CD23 binding affinities ranging from 0.1 nM to 1000 nM, more preferably at least 50 nM, and most preferably at least 5 nM.

Host cells suitable for expression of recombinant immunoglobulins are well known in the art. For example, recombinant antibodies may be expressed in Chinese hamster ovary (CHO) cells, DG44 or DUXB11; or CHO cells CHO K-1; mouse myeloma cells SP2/0 or X63-Ag8.653 or NSO; rat myeloma cells YB2/0; baby hamster kidney cells, BHK; human embryonic kidney line, 293; monkey kidney cells, CV1; human lung fibroblasts, WI38; human cervical carcinoma cells, HELA; insect cells, plant cells, yeast or in bacteria. Further, vectors suitable for expression of immunoglobulins are also well known in the art and are commercially available.

A particularly preferred vector system is the translationally impaired vector system disclosed in <sup>U.S. Patent</sup> ~~U.S. Serial~~ No. 5,448,207 <sup>^</sup> No. 08/147,696 (now allowed), which comprises a translationally impaired dominant selectable marker (neo) containing an intron into which a desired heterologous DNA is inserted. This vector system has been found to provide for very high yields of recombinant proteins, e.g.,

- 21 -

immunoglobulins. However, the subject anti-CD23 antibodies may be produced in any vector system which is suitable for expression of functional immunoglobulins.

Also, the present invention embraces human monoclonal antibodies of the gamma-1 type which are specific to human CD23. Methods for isolation of human monoclonal antibodies are also well known in the art and include *in vitro* methods, e.g., *in vitro* immunization of human B cells in tissue culture, and *in vivo* methods, e.g. synthesis of human monoclonal antibodies in SCID mice. A preferred means of producing human monoclonal antibodies in SCID mice which combines *in vitro* priming of human spleen cells which are then introduced into SCID mice is disclosed in <sup>5,811,524</sup>  
B ~~U.S. Serial No. 08/488,376~~ (incorporated by reference in its entirety herein). This method is advantageous as it provides for the reproducible recovery of monoclonal antibodies having high affinity against a desired antigen, e.g., a human antigen.

Also, the present invention embraces human monoclonal antibodies which compete with the primate anti-human CD23 monoclonal antibodies 5E8 and 6G5 for binding to CD23.

#### EXAMPLE 1

##### Production of Primate Anti-CD23 Antibodies

Five primate monoclonal antibodies specific to CD23 were isolated from macaques substantially according to the



- 22 -

U.S. Patent No. 5,658,570

P methodology disclosed in Serial No. ~~08/379,072~~, which has been incorporated by reference herein. The exact techniques utilized are described in detail below.

**Methodology for Isolation and Characterization  
of Anti-Human CD23 Monoclonal Antibodies**

**Purification of the Immunogen sCD23 from 8866 cells**

During purification, soluble CD23 (sCD23) was quantified by a three-step ELISA using a murine anti-CD23 antibody (Binding Site; catalog # MC112) as a capture. The antigen was partially purified from cultures of 8866 cells maintained in suspension bioreactors using RPMI 1640 (JRH Biosciences; catalog # 56-509) supplemented with 10% fetal bovine serum (JRH Biosciences) and 4 mM glutamine (JRH Biosciences; catalog # 90114) at 37°C. Carbon dioxide was used to maintain pH 7.1. After removing cells by 0.45 µm filtration, phenylmethyl sulfonyl fluoride (final concentration 0.2 mM, Sigman Chemical Co.; catalog # P-7626) and ethylenediaminetetraacetic acid (final concentration 3 mM, Sigma Chemical Co.; catalog # EDS) were added to the supernate and the solution stored at 2-8°C. The cell-free supernate was concentrated approximately 15 to 20-fold using a hollow-fiber ultrafiltration cartridge (A/T Technology; catalog # UFP-10-C-9A; 10,000 d MWCO) or tangential flow ultrafiltration cartridge (Filtron Corporation; 10,000 d MWCO) at ambient temperature. The concentrated supernate was sterile filtered and stored at -70°C.

- 23 -

Thawed concentrates were de-lipidated by adding SM-2 BioBeads (BioRad Industries; catalog # 152-3920) at 5 g/L and stirring overnight at 2-8°C. The resin was removed by filtration and the solution stored at 2-8°C. For some preparations of sCD23, concentrates were fractionated using ammonium sulfate (35-70% (w/v); Fisher; catalog # A702-3) before or after de-lipidation.

The de-lipidated solution was subsequently purified using affinity chromatography at 2-8°C. The affinity matrix was prepared by covalently linking a murine anti-CD23 monoclonal antibody (BU38) to Sepharose using CNBr-activated Sepharose 4B (Sigma Chemical Co.; catalog # C-9142). The BU38 antibody was purified to >90% homogeneity from ascites (Binding Site; catalog # CUS830) using Protein A chromatography. The de-lipidated solution was applied to the affinity column (1.5 x 5 cm) equilibrated with 1XPBS (Gibco BRL; catalog # 70013-0.32), pH 7.2 and the column washed with 1XPBS, pH 7.2, containing 0.05% NP40 (Sigma Chemical Co.) to remove non-bound protein. Soluble CD23 was eluted using 3.5 M MgCl<sub>2</sub> (Fisher; catalog # M33-500). Fractions containing sCD23 were combined and dialyzed (Baxter Spectra/Por; catalog # D1615-1) against 1XPBS, pH 7.2 at 2-8°C. After dialysis, the protein solution was concentrated by centrifugation using Centri-prep 10 spin filters (Amicon Corporation; MWCO 10,000 d) and preparations stored at -70°C. The purity of sCD23 was

estimated to be >70% using SDS-PAGE analysis (4-20% pre-cast gels, Novex Corporation) and Coomassie staining.

**Immunization of Primates and Isolation of Immune Cells**

Cynomolgus monkeys (White Sands Research Center, Alamogordo, New Mexico) were immunized with soluble CD23 which had been purified from the supernatant of human RPMI 8866 cells (B cell lymphoma, Hassner and Saxon, J. Immunol., 132:2844 (1984)). Each monkey was immunized every third week with 200 µg soluble CD23 in 500 µl PBS mixed with 167 µl Temuritide (adjuvant peptide) (Sigma, St. Louis, MO, Catalog # A-9519) and 333 µl 3X PROVAX® (IDEC Pharmaceuticals Corporation). Immunization was effected intradermally, intraperitoneally, intramuscularly and subcutaneously. The titer of anti-CD23 antibodies in the serum of the monkeys was measured by ELISA on 8866 cells and compared to a pre-bleed from the same monkeys.

Monkey PRO 978, with a serum titer of fifty thousand was sacrificed, and the spleen and lymph nodes were surgically removed, and shipped on ice to IDEC pharmaceuticals, submerged in sterile RPMI-1640 (Gibco BRL, Gaithersburg, MD, Catalog # 21870-050) supplemented with 10% fetal calf serum, 2 mM L-glutamine, 2 mM sodium pyruvate and 50 µg/ml gentamicin. Immediately upon arrival the spleen was homogenized by squeezing it through a wire mesh with a glass pistil. Red blood cells were lysed in an ammonium

chloride based hypotonic buffer and the remaining lymphocytes collected and washed in RPMI-1640 at least three times. Lymph nodes were homogenized similarly into a single cell suspension, collected and washed at least three times in RPMI-1640.

#### Production of Hybridomas

After the last wash, the cells were counted, and the primate cells obtained above were then somatically fused to the mouse-human heterohybridoma cell line H6K6/B5 (Carroll et al., *J. Immunol. Methods*, 89:61 (1986)) using standard techniques (Boerner et al., *J. Immunol.*, 147:86 (1991)) and plated into 96 well dishes (175 dishes or 14,700 wells for the spleen, and 17 dishes or 1386 wells for the lymph nodes) at 300,000 cells per well.

This procedure involved the mixing of lymphocytes and the above-identified fusion partner, at a 2:1 ratio, which cells were slowly resuspended into 50% PEG 1500 (Sigma, Catalog # P5402) for 1 minute. These cells were then allowed to rest for 1 minute and then slowly further resuspended in excess RPMI-1640. Afterward, the cells were again allowed to rest, this time for 15 minutes before a light spin at 250 x g. The cells were then resuspended in RPMI-1640 growth media, which was supplemented with 20% Fetal Calf Serum, 2 mM L-Glutamine, Sodium Pyruvate, Non-Essential Amino Acids and 50 µg/ml Gentamicin, con-

- 26 -

taining 100  $\mu$ M Hypoxanthine, 16  $\mu$ M Thymidine (Boehringer-Mannheim, Germany, # 623091) and 5.8  $\mu$ M Azaserine (Sigma, Catalog # A 1164) (HTA). HTA is a selection agent which provides the survival of successfully fused cells (primate lymphocyte fused with heterohybridoma fusion partner).

Approximately 65% of the wells showed growth (10,500 wells). These wells were then screened for the presence of anti-human CD23 antibody by a three step cell ELISA.

#### ELISA Procedure

The first step of the ELISA comprised the transferral of fifty microliters of supernatant from each well to ninety-six well plates which had previously been coated with  $10^5$  8866 cells (CD23 positive cell line) per well. These plates were made by first coating the plates with 50  $\mu$ l of aqueous solution containing twenty  $\mu$ g/ml Poly L-Lysine (Sigma Catalog # P1399, MW 150,000 - 300,000) for thirty minutes at room temperature. The remaining solution was removed ("flicked out") and the plates left to dry. Once dry, fifty  $\mu$ l of 8866 cells in PBS were transferred and spun at 600 g for five minutes. The 8866 cells were covalently bound to the plate by adding fifty  $\mu$ l 0.5% glutaraldehyde (Sigma Catalog # G6257) in phosphate buffered saline (PBS) for 15 minutes. The glutaraldehyde was removed (flicked out) and the plates blocked with one hundred fifty  $\mu$ l 100 mM glycine (Sigma Catalog # G-2879)

- 27 -

in 0.1% BSA - PBS. After the addition of supernatants, the plates were incubated at 37°C for one to two hours and washed seven to nine times with tap water, and a goat anti-human IgG antibody coupled to horse radish peroxidase (HRPO) (Southern Biotech, Birmingham, Alabama, Catalog # 2040-05) diluted 1:2000 into 1% dry skimmed milk (Vons) in PBS - 0.05% Tween 20 (Sigma, Catalog # P1379) was added. The plates were incubated for forty-five minutes at 37°C, and again washed seven to nine times in tap water. The presence of the HRPO was detected by a color development after the addition of a TMB reagent (Kirkegaard & Perry, Gaithersburg, MD, Catalog # 50-76-02 and 50-65-02), 100  $\mu$ l/well. The reaction was stopped by adding twenty-five  $\mu$ l 4N H<sub>2</sub>SO<sub>4</sub>. Optical density (OD) was measured at 470 nm on a spectrophotometer (Titertek Multiscan). The OD values greater than two times the background were scored as positive.

The second step in the ELISA was effected to confirm that the supernatants which had been scored positive in the first ELISA reacted to CD23 and not to some irrelevant antigen. This was effected by testing the supernatants on Supt1 cells (ERC BioServices Corporation, Rockville, MD, Catalog # 100), a CD23 negative human cell line, using the same ELISA procedure. Supernatants that scored similarly in both tests were discarded. These results indicated that fifty-six of the 10,500 wells with growth showed the

- 28 -

presence of a primate monoclonal antibody that bound to 8866 cells in two separate screenings at different times and did not bind to SupT1 cells.

The third step of the ELISA was conducted to determine whether the supernatants identified according to the first two ELISA steps, reacted with soluble CD23. In this third ELISA, 96 well plates were coated at 4°C overnight with 2 µg/ml BG-6 (Biosource International, Camarillo, CA, Catalog # CT-CD23-CF), a mouse monoclonal antibody that binds to soluble CD23 but does not block CD23-IgE binding, contained in a 50 mM bicarbonate buffer, pH 9.3. After removing the coating buffer, fifty µl of semi-purified soluble CD23 at a predetermined dilution in PBS were added to the plate and incubated for two hours at room temperature. After washing the plate with tap water seven to nine times, 50 µl supernatants from selected wells were added. After washing the plate in tap water seven to nine times, 50 µl rabbit anti-human IgG (mouse adsorbed)-HRPO (Southern Biotech, Catalog # 6145-05) diluted 1:4000 in 1% dry skimmed milk in PBS with 0.05% Tween 20 were added, incubated for two hours at 37°C, washed seven to nine times in tap water and developed with TMB as described above. Wells with OD's greater than two times the background were again scored as positive.

Twenty-one of the fifty-six wells that showed binding to 8866 cells also bound to sCD23 in the ELISA. These

wells were expanded and subcloned at least twice by plating out cells at one cell per three wells. After approximately three months, five stable hybridomas producing primate monoclonal antibodies to CD23 were obtained.

#### Antibody Purification by Protein A Methods

Essentially, antibodies are purified by centrifugation of the culture supernatant to remove cells and debris. The resultant centrifuged samples are then filtered through a 0.2 $\mu$ m filter. A protein A sepharose Fast flow column is then prepared and equilibrated using PBS (pH 7.4). The supernatant is then loaded on the column at an appropriate flow rate (2 ml/min). After loading, the wash column is washed with 10 column volume of PBS (pH 7.4). The antibody is then eluted from the column with elution buffer (0.2 M acetic acid, 0.1 M glycine pH 3.5) at 1 ml/min flow rate. One milliliter fractions/tube (2.0 M Tris-Hcl pH 10.0) including 100 $\mu$ l of Tris, are then collected. Afterward, spectrophotometer readings are taken at 280 nm. The resultant fractions with high absorbance at 280 nm containing the antibody are then collected and dialyzed against PBS overnight. The product is then sterilized by filtration through 0.22 $\mu$ m membrane and stored at -20°C.

Four of these five primate anti-human CD23 monoclonal antibodies (1H6, 2C8, 5E8 and 6G5) were demonstrated to



- 30 -

inhibit IgE production in an *in vitro* assay which measures IgE production by IL4-hydrocortisone induced peripheral blood mononuclear cell (PBMC) cultures. These results are shown in Figure 1. The assay conditions are described below. The fifth primate monoclonal anti-human CD23 antibody B3B11 was inactive in this assay.

IL-4 Stimulated IgE Production  
by Peripheral Blood Mononuclear Cells

As discussed supra, the subject primate antibodies and PRIMATIZED® forms thereof were assessed for their ability to inhibit IgE production in an *in vitro* assay which measured the effect of such antibodies on IgE production by IL-4 stimulated peripheral blood mononuclear cells.

Materials for in vitro IL-4 IgE Assay

Forty-eight well flat bottom cluster plates (Costar Catalog # 3548) (1.5 million PBMCs per ml per well (48 well plate))

Human recombinant IL-4 (Genzyme Catalog # 2181-01; 10 µg (2.5 x 10<sup>7</sup> units).

anti-CD23 Mabs:

murine Mab (MHM6; DAKO. Catalog # M763)

primate Mabs (no preservatives)

PRIMATIZED® (no preservatives)

- 31 -

HB 101 basal medium: (Irvine Scientific Catalog # T000)

HB101 supplement: (Irvine Scientific Catalog # T151)

Fetal Bovine Serum: (FBS; Bio-Whittaker Catalog # 14-501F)

dimethylsulfoxide: (DMSO; Fisher Scientific Catalog # D128-500)

hydrocortisone: (Sigma Catalog # H-0888)

puromycin: (Sigma Catalog # P-7255)

cyclohexamide: (Sigma Catalog # C-7698)

*B* HISTOPAQUE  
HISTOPAQUE®: (Sigma Catalog # H-8889)

Hank's Buffered Salt Solution: (HBSS; Irvine Scientific Catalog # 9232)

1% FBS in HBSS

concentrated Dulbecco's phosphate buffered saline (10x DPBS; Bio-Whittaker, Catalog # 17-517Q)

Bath Clear Microbicide (Fisher, Catalog # 13-641-334) in DPBS

Solutions:

puromycin solution: 40 µg/ml in HB101 growth medium

cyclohexamide solution: 200 µg/ml in HB101 growth medium

hydrocortisone solution: 0.1 M solution in DMSO

anti-CD23 murine Mabs were extensively dialyzed to remove preservatives

## HB101 growth medium

HB101 basal medium	500 ml
HB101 supplement in 10 ml	
sterile filtered distilled H <sub>2</sub> O	5 ml
FBS	10 ml
hydrocortisone solution	
(final conc. 5 $\mu$ M)	0.25 ml

In Vitro Assay Procedure

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2

using a hemocytometer. Typically, 300 to 600 million PBMCs can be recovered from a single 450 ml buffy coat package.

The collected PBMCs are then washed three times in 1% FBS/HBSS. The washed cells are collected by centrifugation for seven minutes at 1300 rpm at 7°C.

The number of cells collected is then determined using a hemocytometer. The cell concentration is adjusted to about three million cells per milliliter of HB101 growth medium.

Approximately about 1.5 million cells (0.5 ml) are then added to each well of a 48 well plate. In general, five replicate samples are prepared for each experiment. The perimeter wells of each plate are not used for cell samples. Accordingly, these wells are filled, e.g., using 0.5 ml of 0.05% BathClear/DPBS.

0.5 ml HB101 growth medium containing desired amounts of IL-4 and Mab is then added to the wells. The IL-4 used is recombinant DNA-generated human interleukin 4. The Mab used in the assay is a murine, primate or PRIMATIZED® antibody. Typically, IL-4 is added at a final concentrate of 100 U/ml and Mab is added at a final concentrate ranging from 0.01 to 3 µg/ml.

The cells are then incubated for nine to eleven days at 37°C in a moist incubator set at 5% CO<sub>2</sub>. After incuba-

tion, the supernatant fluids are collected and the IgE content is measured.

### IgE ELISA

The following list identifies materials and solutions used in the IgE ELISAs.

#### Materials and Solutions Needed for IgE ELISA

sulfuric acid, 4 M

coating buffer: 10 mM sodium bicarbonate buffer, pH 9.6

concentrated phosphate buffered saline (10x PBS) stock solution:

NaH <sub>2</sub> PO <sub>4</sub>	26.6 gm
Na <sub>2</sub> HPO <sub>4</sub>	289 gm
NaCl	1064 gm
distilled H <sub>2</sub> O	10 L

blocking buffer: 10% FBS/PBS

dilution buffer: 1% BSA/0.05% Tween 20/PBS

washing buffer: 0.05% Tween 20/PBS

goat anti-human IgE (epsilon chain-specific), unlabeled:  
(Tago Catalog # 4104)

human IgE standard: (The Binding Site Catalog # BP094)

goat anti-human IgE, HRP-labeled: (Tago Catalog # AHI  
0504)

TMB peroxidase substrate: (KPL Catalog # 50-76-02)

peroxidase solution B: (KPL Catalog # 50-65-02)

working substrate solution: mix substrate and Solution B at 1:1 ratio

Immulon II microtiter plates (Dynatech Labs Catalog # 011-010-3455)

IgE ELISA Procedure

Each well of a microtiter plate is coated using 100  $\mu$ l of a coating buffer containing 2  $\mu$ g/ml goat anti-human IgE.

The coated plate is then incubated overnight at 4°C.

After incubation, each well in the plate is then washed three times with 200  $\mu$ l of Tween 20/PBS. After washing, the non-specific binding sites are blocked with 200  $\mu$ l blocking buffer/well for 1 hour at 37°C.

One hundred  $\mu$ l of samples or standards are then added to each well; which wells are then incubated overnight at 4°C. After incubation, the samples are tested with or without dilution. A standard concentration curve is prepared for each plate using several dilutions of IgE ranging from 0.1 to 50 ng/ml.

After overnight incubation, each plate is washed five times with Tween 20/PBS.

One hundred  $\mu$ l of horseradish peroxidase (HRP) labeled goat anti-human IgE diluted 1:10,000 in dilution buffer is then added to each drained well. The plate is then incubated for 4 hours at 37°C.

The plates are then washed 5 times with Tween 20/PBS and 3 times with water.

One hundred  $\mu$ l of 3,3',5,5'-tetramethylbenzidine working substrate solution is then added to each well. The plate is then incubated for twenty-five minutes in the dark at room temperature. After incubation the developing reaction is stopped by the addition of fifty  $\mu$ l of 4 M sulfuric acid.

The absorbency is then read concurrently at 450 and 540 nm. The 540 nm absorbency values are subtracted as background.

Assay for  $K_d$  measurement of primate monoclonal anti-human CD23 antibodies

Scatchard Analysis Procedure

1. Radiolabeling Procedure

iodo-BEADS are washed with 100mM Phosphate Buffer, pH 7.4 twice using 1 mL of buffer per 2 beads. The beads are then dried on filter paper.

The two beads are then added to 100  $\mu$ l  $^{125}$ I solution, containing about 1mCi of I, diluted with 200  $\mu$ l of the phosphate buffer, and left at room temperature for 5 minutes.

The antibody (50  $\mu$ gs) is added to the preloaded beads. The reaction time for maximal incorporation of radioactivity is 6 minutes.

The reaction is stopped by removing the radiolabeled antibody from the reaction vessel.

- 37 -

Gel filtration is then performed to remove excess  $^{125}\text{I}$  or unincorporated  $^{125}\text{I}$  from the radiolabeled antibody solution. This is effected by passing the radiolabeled antibody over a column made up of 1.5mL Sephadex-G25, 1.5 mL DEAE Sephadex-A25 and 0.5mL Amberlite. The radiolabeled antibody is eluted off in a total volume of 5 mL at a concentration of about 10  $\mu\text{g/mL}$ . (Elution Buffer: 1XPBS containing 10% Gelatin, 2% Sodium Azide and 1% BSA).

## 2. Optimization Assay (Direct Binding Study)

The specific activity of the 10  $\mu\text{g/mL}$  radiolabeled solution is determined by taking a 1  $\mu\text{l}$  sample and running the sample on a gamma counter.

### Example:

- $1 \times 10^5 \text{ cpm}/\mu\text{l} \times 1000 \mu\text{l}/10\mu\text{g antibody}$   
 $1 \times 10^5 \text{ cpm}/\mu\text{g antibody}$   
 $1 \times 10^4 \text{ cpm}/\text{ng antibody}$
- Molecular wt. of antibody = 75,000 ng/nmole
- **Specific activity:**  
 $1 \times 10^4 \text{ cpm}/\text{ng} \times 75,000 \text{ ng}/\text{nmole} = 7.5 \times 10^8 \text{ cpm}/\text{nmole}$

The antigen-coated plate is blocked (to eliminate non-specific binding, e.g., with mB7.1-CHO) and the background plate (i.e., Untransfected-CHO) for one hour at room temperature-with 200  $\mu\text{l}/\text{well}$  of blocking buffer



(Blocking Buffer: 1xPBS containing 10% Gelatin, 2% Sodium Azide, 1% BSA and 10% FBS).

The plate(s) are then washed, typically ten times by hand with tap water.

The 10  $\mu\text{g/mL}$  radiolabeled antibody (50  $\mu\text{ls}$ ) is then titrated by two-fold serial dilutions across the plate(s) using a multichannel pipette. Incubate for one hour at room temperature.

The plate(s) are again washed about 6-7 times with 200  $\mu\text{l/well}$  of wash buffer (Wash Buffer:--1xPBS containing 10% Gelatin and 2% Sodium Azide).

The radio activity counts in each well are then determined by running the wells on a gamma counter.

The optimal radiolabeled antibody concentration is the concentration in which the difference between the specific counts and background counts is at a maximum.

### 3. Scatchard Analysis of Competition Assay

The 10  $\mu\text{g/mL}$  radiolabeled solution is diluted to the optimal concentration determined in the **Direct Binding experiment**.

The antigen-coated plate and the background plate are blocked for one hour at room temperature with 200  $\mu\text{l/well}$  of blocking buffer.

The plate(s) are then washed, e.g., about 10 times, by hand with tap water.

The "cold" (no radiolabel) antibody is then titrated by two-fold dilutions in a separate U-bottom microtitre plate. The starting concentration of the "cold" antibody should be at least 100 times greater than that of the optimal radiolabeled antibody concentration.

**Example:**

- Optimal Radiolabeled Conc.: 0.5  $\mu\text{g/mL}$
- "Cold" Antibody Conc.: 100  $\mu\text{g/mL}$  (Note: 1:2 titration in the first well will adjust the "cold" antibody concentration to 50  $\mu\text{g/mL}$ .)

Fifty  $\mu\text{l}$ /well of optimal radiolabeled antibody are then added to the wells containing "cold" antibody.

One hundred  $\mu\text{l}$ /well of the mixed solution are then transferred to the corresponding wells of the antigen-coated plate, and incubated for one hour at room temperature.

Also, it is desirable also that the following controls be effected:

- a) Direct binding of radiolabeled antibody to antigen-coated plate (5 wells),
- b) Direct binding of radiolabeled antibody to background plate (5 wells).

After incubation, the plate(s) are washed, e.g., about 6-7 times, with 200  $\mu\text{l}$ /well of wash buffer.

- 40 -

The radio activity counts in each well are then determined by running the wells on a gamma counter.

These calculations are determined by calculating the specific counts in each well tested by subtracting the background counts from the counts bound to the antigen-coated plate.

#### 4. Calculations for Scatchard Analysis

The Molar Concentration of Bound antibody [B] can then be determined as follows:

**Example:** At 50  $\mu\text{g/mL}$  "cold antibody"

Specific counts bound: 4382 cpm

Counts bound in the presence of 50  $\mu\text{g/mL}$  "cold" ab: 215 cpm

Difference: 4382 cpm - 215 cpm = 4167 cpm

Specific Activity (radiolabeled ab):  $5.54 \times 10^9$  cpm/nmole

$4167 \text{ cpm} + 5.54 \times 10^9 \text{ cpm/nmole} = 7.52 \times 10^{-7} \text{ nmole}$

$7.53 \times 10^{-7} \text{ nmole} + 0.05 \text{ mL (sample vol.)}$

$$= 1.50 \times 10^{-5} \text{ nmole/mL}$$

$$= 1.50 \times 10^{-8} \text{ } \mu\text{mole/mL}$$

$$[\text{B}] = 1.50 \times 10^{-11} \text{ mole/mL (M)}$$

**Total Molar Concentration [T]** is determined as follows:

$$50 \text{ } \mu\text{g/mL} \times 1 \text{ } \mu\text{mole/75,000 } \mu\text{g} = 6.67 \times 10^{-4} \text{ } \mu\text{mole/mL}$$

$$= 6.67 \times 10^{-7} \text{ mmole/mL (M)}$$

$$[\text{T}] = 66667 \times 10^{-11} \text{ mmole/mL (M)}$$

- 41 -

Free antibody [F] is determined as follows:

Free Molar Conc. = Total minus Bound

$$\begin{aligned} [F] &= (66667 \times 10^{-11}) - (1.50 \times 10^{-11}) \\ &= 66665.5 \times 10^{-11} \text{ mmole/mL (M)} \end{aligned}$$

Calculate B/F.

Plot B versus B/F on <sup>CRICKET GRAPH</sup>~~Cricket-Graph~~ software.

Activity and Affinity of Anti-Human CD23  
Antibodies According to the Invention

Four of the five isolated primate anti-human CD23 monoclonal antibodies (B3B11, 2C8, 5E8 and 6G5) were found to inhibit IgE production in the above-identified in vitro assay which measures IgE production by IL4-hydrocortisone induced peripheral blood mononuclear cell (PBMC) cultures. These results are shown in Figure 1. The fifth primate monoclonal anti-human CD23 antibody 3G12 was inactive in this assay.

Two of the four primate monoclonal anti-human CD23 antibodies (B3B11 and 2C8) found to be active in this in vitro assay were found to compete with a commercially available mouse anti-human CD23 antibody MHM6 (CAKO A/S, Glostrup, Denmark Catalog # M763). (Figure 2, top panel.) However, in repeated assays these antibodies were not as potent IgE inhibitors as MHM6 (data not shown). By contrast, the other primate anti-CD23 monoclonal antibodies (5E8 and 6G5) were found to compete with each other and did not compete with MHM6. (Figure 2, middle and bottom

panels.) Moreover, the primate anti-human CD23 monoclonal antibody 5E8 was found to be a potent inhibitor of IL-4 induced IgE in the *in vitro* assay. (See Figures 1 and 3.)

Modified Hu-SCID-Mouse Model for Human IgE Synthesis  
and Measuring the Inhibition of IL-4 Induced IgE  
Production by Anti-CD23 Antibodies In Vivo

A modified hu-PBMC-SCID mouse model was also developed to detect the effect of the subject antibodies on induced human IgE production *in vivo*. PBMCs obtained from two donors were cultured with IL-4 *in vitro* for two days. PBMCs were pooled and used to reconstitute groups of C.B.-17 SCID mice with and without antibodies. Mice were bled on day 14, 21, 28 and 35 and serum IgG and IgE levels were determined by ELISA. This *in vivo* model was used to assay primate and two different versions of PRIMATIZED® antibodies to CD23 for their ability to inhibit the production of IgE.

A modified SCID mouse model was used because it is known that severe combined immunodeficiency scid/scid (SCID) mice, C.B.-17 (Bosma et al., *Nature*, 301:527 (1983)) reconstituted with human peripheral blood mononuclear cells (hu-PBMC-SCID) can produce significant quantities of human immunoglobulins (Ig) (Moiser et al., *Nature*, 335:256 (1988); Moiser et al., *J. Clin. Immunol.*, 10:185 (1990); Abedi et al., *J. Immunol.*, 22:823 (1992); and Mazingue et al., *Eur. J. Immunol.*, 21:1763 (1991).) The

predominant isotype of human immunoglobulin (Ig) produced in hu-PBMC-SCID mice is IgG. Generally, IgM, IgA and IgE isotypes are found in very low or non-detectable levels except in cases where PBMC is obtained from donors with certain autoimmune or allergic disease conditions. It has also been reported that manipulation of hu-PBMC SCID mouse model with certain cytokines may be provided for the generation of significant levels of non-IgG isotypes, including IgE (Kilchherr et al., *Cellular Immunology*, 151:241 (1993); Spiegelberg et al., *J. Clin. Investigation*, 93:711 (1994); and Carballido et al., *J. Immunol.*, 155:4162 (1995)). The hu-PBMC-SCIDs, has been also used to generate antigen specific Ig provided the donor has been primed for the antigen *in vivo*.

Therefore, the aim of the present inventors was focused on establishing a suitable human IgE producing hu-PBMC-SCID mouse model that could be used to test the efficacy of therapeutic for treatment of IgE related diseases such as allergic disorders, including the subject anti-CD23 antibodies.

#### **Materials and Methods:**

The following materials and methods were used in the hu-PBMC-SCID mouse model described below.

**SCID mice:** C.B-17 scid/scid immunodeficient mice were obtained from Taconic (C.B. -17/IcrTac-scid<sup>EDF</sup>) and main-

tained in IDEC Pharmaceuticals' animal facility. Mice were housed in sterilized microbarrier units with sterilized bedding. Animal studies were performed in accordance with the "Guide for the Care and Use of Laboratory Animals" specified by the Committee on Care of Laboratory Animal Resources Commission on Life Science-National Research Council (Guide for the Care and Use of Laboratory Animals, DHHS Publ. No. (NIH) 86-23, Bethesda, MD, NIH, 1985).

**Human PBMC:** PBMCs were isolated from buffy coats obtained from a blood bank by centrifugation through Ficoll-Hypaque (Histopaque-1077) as recommended by the manufacturer (Sigma Diagnostics Catalog # 1077-1). Lymphocyte preparation at the interface of the gradient were harvested and washed three times in Hanks Balanced Salt Solution (HBSS) (Bio-Whittaker Catalog # 10-527F). For each experiment PBMCs were obtained from two separate donors and cultured separately *in vitro*. PBMCs were resuspended at  $1-3 \times 10^6$  cells/ml concentration in HB-Basal medium plus 1% HB101 lyophilized supplement (Irvine Scientific Catalog # T000 & T151) containing 5% FCS plus 1000 IU/ml of IL-4 (Genzyme, Inc. Catalog # 2181-01) and incubated for 48 hours at 37°C with 5% CO<sub>2</sub>. After incubation, the cells from different buffy coats were harvested, pooled and used to reconstitute SCID mice.

In Vivo Assay Conditions

Groups of mice (four to five per group) were injected with fifty-sixty x  $10^6$  lymphocytes in 200-300  $\mu$ l volume of HBSS intraperitoneally (i.p.) on day zero. For the groups that received anti-CD23 antibody, on day zero, PBMCs were mixed with anti-CD23 antibody (200 to 400  $\mu$ g/mouse) before i.p., injection and the second injection was given on day seven. All mice received 5000 IU per mouse of IL-4 i.p., between day zero to day five. A group which was not injected with antibody served as the control group. Mice were bled from a retro-orbital vein and the serum was analyzed for IgG and IgE on days fourteen, twenty-one, twenty-eight and thirty-five by ELISA.

Figure 8 shows that the primate anti-human CD23 monoclonal antibody 5E8 is effective in inhibiting IL-4 induced IgE production in vivo in the SCID mouse model.

Cloning and Expression of PRIMATIZED® ( anti-human CD23 Monoclonal Antibodies

In order to clone primate immunoglobulin variable domains, Poly A+ RNA was separately isolated from approximately  $2 \times 10^6$  cells from the primate heterohybridomas secreting the anti-human CD23 monoclonal antibodies 6G5 and 5E8 by using the Micro-FastTrack mRNA isolation Kit (Invitrogen Catalog # K1520-02) according to methods set forth by the manufacturer.



The first strand of cDNA was synthesized from the poly A+ RNA by using the cDNA Cycle Kit (Invitrogen Catalog # L1310-01) according to conventional methods.

The light and heavy chain variable regions of 6G5 and 5E8 were then isolated by PCR from cDNA using PCR primers that were selected based upon different consensus families of human immunoglobulins. 5' primers were selected which corresponded to the beginning of the leader sequences of the light and heavy variable region and 3' primers were selected which corresponded to the J region (The specific primers used to PCR amplify the lambda light chain variable domain of 6G5, the kappa light chain variable domains of 5E8, and the heavy chain variable domains of 6G5 and 5E8 are set forth in Tables 1-3). PCR was performed according to standard methods (30 cycles with 1 minute at 94°C, 1.5 minutes at 54°C and 2 minutes at 72°C in a Hot start 100 tube (Gibco BRL Catalog # 10332-013). PCR was set up in 50µl reactions containing 5µl out of 80µl cDNA (from 2x10<sup>6</sup> cells) as a template, 2µl of 5 nM dNTP, 1µl of Taq polymerase, 5µl of Taq polymerase buffer, 2µl of the 5' primer (25 pmoles/µl), 2µl of the 3' primer (25 pmoles/µl), and 36µl of water. (Taq polymerase and buffer were obtained from Stratagene Catalog # 600131, dNTP from Boehringer Mannheim Catalog #1581295.)

A) Construction of the plasmids N5LG1 + 6G5 and N5LG4P + 6G5

1) Cloning the light chain variable domain of primate monoclonal anti-human CD23 antibody 6G5 by PCR

B The first PCR amplification of the light chain variable region from the cDNA of primate monoclonal antibody 6G5 showed bands which were consistent in situ with the lambda light chain variable region. These bands appeared in all reactions using the three different early leader [SEQ ID NOS.: 5--21] sequence primers. (See Tables 1-3.) However, the PCR product obtained using primer 745 (Family 2) was considered more specific because of the relatively greater intensity of the PCR product band.

This PCR product was isolated using a Qiaquick Gel Extraction Kit (Qiagen Catalog # 28704). The purified PCR fragment was digested with Bgl II and Avr II restriction endonucleases, and ligated into the mammalian expression vector N5LG1 which was digested with the same restriction endonucleases. Twenty microliters of the ligation mixture containing the purified PCR product from one fifty microliter PCR reaction, 100 mg N5LG1 vector, two microliters of 10x ligation buffer (NEB Catalog # 202S) and two microliters of T4 ligase (NEB Catalog # 202S), were then incubated at 14°C overnight.

The mammalian expression vector N5LG1 contains genetic sequences (e.g., regulatory sequences, coding sequenc-

es) which provide for the expression of four separate proteins in a mammalian cell. They are:

(i) a partial immunoglobulin light chain with the human lambda light chain constant region and unique restriction endonuclease sites for inserting light chain variable domains;

(ii) a partial immunoglobulin heavy chain with the and human gamma 1 chain constant region coding sequences and unique restriction endonuclease sites for inserting heavy chain variable domains;

(iii) a neomycin phosphotransferase gene used to select for cells that have incorporated the plasmid and are resistant to the antibiotic Geneticin (Gibco BRL Catalog # 10131-1209); and

(iv) a murine dihydrofolate reductase gene (DHFR) which provides for the selection and genomic amplification when cells are cultured in the presence of methotrexate (MTX, Sigma Catalog # A-6770) (Reff et al., *Blood*, 83:433-445 (1994)).

After ligation, the mixture was digested using Pme I restriction endonuclease, which digests the parent N5LG1 plasmid, but not the N5LG1 plasmid which has been ligated to the light variable domain of 6G5. After digestion, the mixture was transformed into <sup>EPICURIAN COLI XL1-BLUE</sup> ~~Episurian coli~~ <sup>XL1-Blue</sup> competent cells (Stratagene Catalog # 200249) as follows.

- 49 -

One hundred microliters of competent cells were mixed with 10  $\mu$ l of the above ligation mixture, set on ice for 30 minutes, then heated at 45°C for 30 seconds. This mixture was placed on ice for 2 minutes, and 900  $\mu$ l of SOC, prewarmed to room temperature, was then added. (SOC is LB broth Gibco BRL Catalog # 10855-013, plus 0.02 M  $MgCl_2$ , 0.02 M  $MgSO_4$ , and 0.02 M D-glucose.) After incubation at 37°C for an hour, the mixture was centrifuged at 4000 g for a minute, and 800  $\mu$ l of supernatant discarded. The rest of the mixture was plated onto a LB agar (Gibco BRL Catalog # 12945-044) dish containing 50  $\mu$ g/ml ampicillin (Amp, Gibco BRL Catalog # 13075-015). Plasmid DNA was isolated from individual colonies of *E. coli* that grew on the Amp plate by using the <sup>WIZARD</sup> Wizard® Miniprep DNA purification system (Promega Catalog # A7510).

The isolated plasmid DNA was then characterized by digestion with Bgl II and Avr II followed by agarose gel electrophoresis. An ethidium stained DNA band of 400 bp was indicative of a potential successful cloning of a light chain variable domain.

To confirm this was an immunoglobulin light chain variable domain, sequencing was done using the Sequenase 7-Deaza -dGTP DNA Sequencing Kit (USB catalog # 70990) with sequencing primers 607 and GE 108. (See Sequencing [SEQ ID NOS: 22-31] Primers in Table 4.)

- 50 -

A second independent PCR amplification of the light chain from cDNA of primate monoclonal antibody 6G5 was effected using a 5' primer early leader sequence of lambda light chain family 2 (primer 745) and the 3' J region primer 926. (See Primers for PCR of the lambda light chain variable domain of 6G5 in Tables 1-3.)<sup>[SEQ ID NOS.: 5-21]</sup> The isolated PCR product (see technique above) was cloned into TA vector by using the Original TA Cloning Kit (Invitrogen Catalog # K2000-01). The isolated miniprep DNA (see technique above) was examined under agarose gel electrophoresis after digestion with EcoR I restriction endonuclease. The resultant PCR product comprised in the TA vector was then sequenced (as described previously) using Sp6 and M13(-40) forward primers (See Sequencing primers in Table 4).<sup>[SEQ ID NOS.: 22-31]</sup> The resultant light chain sequence was identical to that of light chain from the first PCR. This entire sequence<sup>[SEQ ID NO.: 1]</sup> of the light chain variable domain of primate monoclonal anti-human CD23 antibody 6G5 is presented below.

Light chain variable region of primate monoclonal antibody anti-human CD23 6G5 Leader

Met Ala Trp Thr Leu Leu Leu Val Thr Leu Leu Thr Gln Gly Thr  
 ATG GCC TGG ACT CTG CTC CTC GTC ACC CTC CTC ACT CAG GGC ACA

-1

Gly Ser Trp Ala  
 GGA TCC TGG GCT

## Framework 1

CDR 1

## Framework 2

CDR2

### Framework 3

57                      60    70

Gly Val Ser Asp Arg Phe Ser Gly Ser Lys Ser Gly Asn Thr Ala

GGG GTC TCT GAT CGC TTC TCT GGC TCC AAG TCT GGC AAC ACG GCC

- 52 -

80

Ser Leu Thr Ile Ser Gly Leu Gln Ala Glu Asp Glu Ala Asp Tyr  
TCC CTG ACC ATC TCT GGG CTC CAG GCT GAG GAC GAG GCT GAT TAT

88

Tyr Cys  
TAC TGT

CDR 3

89 90 95 95A 96 97  
Cys Ser Tyr Thr Thr Ser Ser Thr Leu Leu  
TGT TCA TAT ACA ACC AGT AGC ACT TTG TTA

Framework 4

98 100 106 106A 107  
Phe Gly Arg Gly Thr Arg Leu Thr Val Leu Gly  
TTC GGA AGA GGG ACC CGG TTG ACC GTC CTA GGT

2) Cloning the heavy chain variable domain of primate monoclonal anti-human CD23 antibody 6G5 by PCR

The first PCR amplification of the heavy chain variable domain from cDNA of primate monoclonal antibody 6G5 was performed by using the set of early leader sequence primers described supra and the 3' J region primer GE244. <sup>[Seq ID No: 20-31]</sup>  
These primers are in Tables 1-3 <sub>infra</sub>. This reaction resulted in a 350 base PCR product. This 350 base product (purified as described supra), was digested with Nhe I and Sal I, and ligated into N5LG1 and digested with the same endonucleases in the first PCR amplification. The resultant ligation mixture was transformed into host cells

- 53 -

using the same techniques for cloning the light chain. Plasmid N5LG1 containing the 350 base PCR product was then isolated and sequenced (using sequencing primers 266 and 268). (These Sequencing primers are set forth in Table

B  
4.) [SEQ ID NOS: 22-31]  
A

Sequencing revealed that the PCR product contained only part of the heavy variable domain and comprised a deletion in its amino terminus (Sequence began at framework 2, codon 36).

A second independent PCR reaction was conducted to amplify and isolate the heavy chain variable domain of primate monoclonal antibody 6G5 using a 5' early leader sequence primer for family 1 (MB1503) and a 3' J' region primer GE244. (These primers are also contained in Tables

B  
1-3.) [SEQ ID NOS: 5-21]  
A

The resultant PCR product was then cloned into the N5LG1 using the same techniques described supra. Its sequence was found to be identical to the first PCR product.

Therefore, in order to clone the whole heavy variable domain of 6G5 including the missing 5' terminus a new longer 3' primer (MB1533) which included the CDR3 and framework 4 regions of the 6G5 heavy variable chain was then used in a third independent PCR reaction with the family 1 5' primer (MB1503). (These primers are also

B  
contained in Tables 1-3.) [SEQ ID NOS: 5-21]  
A



- 54 -

After PCR, a larger 420 base PCR product was observed on the agarose gel. This PCR product was isolated as described previously, and cloned into a TA vector. The resultant PCR product contained in the TA vector was then sequenced. Sequencing revealed that this DNA contained the whole heavy variable domain and that the 3' part was identical to that of previously cloned partial heavy chain variable domain from the first two PCR reactions.

A fourth independent PCR was performed using the same primers as the third PCR amplification. This resulted in a PCR product which was isolated and cloned into the TA vector as described previously. The sequence of the fourth independent PCR product was found to be identical to that obtained in the third PCR amplification. This sequence, <sup>[Seq ID NO.: 2]</sup> which comprises the heavy chain variable domain of primate monoclonal anti-human CD23 antibody 6G5, is presented below.

**Heavy chain variable region of primate monoclonal antibody anti-human CD23 6G5**

**Leader**

Met Lys His Leu Trp Phe Phe Leu Leu Leu Val Ala Ala Pro Arg  
ATG AAA CAC CTG TGG TTC TTC CTC CTC CTG GTG GCA GCT CCC AGA

-1

Trp Val Leu Ser  
TGG GTC CTG TCC

- 55 -

Mature Protein (Numbering is Kabat)

Framework 1

1		10
Gln	Leu	Gln
Leu	Gln	Glu
Ser	Gly	Pro
Gly	Val	Val
Lys	Pro	Ser
CAG CTG CAG CTG CAG GAG TCG GGC CCA GGA GTG GTG AAG CCT TCG		
20		30
Glu	Thr	Leu
Ser	Leu	Thr
Cys	Ala	Val
Ser	Gly	Gly
Ser	Val	Ser
GAG ACC CTG TCC CTC ACC TGC GCT GTC TCT GGT GGC TCT GTC AGC		

CDR 1

31		35	35a
Ser	Ser	Asn	Trp
Trp	Trp	Thr	
AGT AGT AAC TGG TGG ACC			

Framework 2

36		40		49
Trp	Ile	Arg	Gln	Pro
Pro	Gly	Lys	Gly	Leu
Glu	Trp	Ile	Gly	
TGG ATC CGC CAG CCC CCA GGG AAG GGA CTG GAG TGG ATT GGA				

CDR2

50		52	52A	53		60
Arg	Ile	Ser	Gly	Ser	Gly	Gly
Ala	Thr	Asn	Tyr	Asn	Pro	Ser
Leu						
CGT ATC TCT GGT AGT GGT GGG GCC ACC AAC TAC AAC CCG TCC CTC						
65						
Lys	Ser					
AAG AGT						

```

66              70              80
Arg Val Ile Ile Ser Gln Asp Thr Ser Lys Asn Gln Phe Ser Leu
CGA GTC ATC ATT TCA CAA GAC ACG TCC AAG AAC CAG TTC TCC CTG
      82 82a 82b 82c 83              90
Asn Leu Asn Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys
AAC CTG AAC TCT GTG ACC GCC GCG GAC ACG GCC GTG TAT TAC TGT
      94
Ala Arg
GCC AGA

```

95	100	100a	100b	100c	100d	101	102
Asp Trp Ala Gln Ile Ala Gly Thr Thr Leu Gly Phe							
GAT TGG GCC CAA ATA GCT GGA ACA ACG CTA GGC TTC							

103						110			113	
Trp	Gly	Gln	Gly	Val	Leu	Val	Thr	Val	Ser	Ser
TGG	GGC	CAG	GGA	GTC	CTG	GTC	ACC	GTC	TCC	TCA

In order to insert the cloned heavy chain variable domains of 6G5 into a mammalian expression vector, the heavy chain variable domain in the TA vector (obtained in the 3rd independent PCR) was digested with Nhe I and Sal I and cloned into the N5LG1 vector which was digested with the same restriction enzymes and which vector already contains the light chain variable domain. The resultant mammalian expression vector was named N5LG1 + 6G5.

- 57 -

To construct the N5LG4P + 6G5 vector, both the light and heavy chain variable domains were isolated from N5LG1 + 6G5 by digestion of Bgl II and Avr II, and Nhe I and Sal I respectively. The mammalian expression vector N5LG4P vector is identical to the N5LG1 vector described above, except the human gamma 1 was replaced with a human gamma 4 constant region containing a mutation of a serine to a proline in the hinge region to increase stability of the immunoglobulin and improve pharmacokinetics *in vivo* ("P" mutation). The light chain variable domain was cloned in the plasmid first and the heavy chain variable domain was cloned into the vector containing the light chain variable domain using techniques previously described. This mammalian expression vector was named N5LG4P + 6G5.

**B. Construction of the plasmids N5KG4P + 5E8, N5KG1 + 5E8, N5KG4P + 5E8N-, and N5KG1 + 5E8N-**

**1. Cloning the light chain variable domain of primate monoclonal anti-human CD23 antibody 5E8 by PCR**

The first PCR reaction of the light chain variable domain from 5E8 cDNA was carried out using a set of kappa early leader sequence primers and the 3' J region primer GE204. (See primers for PCR of the kappa light chain variable domain of 5E8 in Tables 1-3). <sup>[SEQ ID Nos.: 5-27]</sup> A 420 base PCR product was obtained. The isolated 420 base PCR product was digested with Bgl II and BsiW I restriction endo-

nucleases, cloned into the mammalian expression vector N5KG4P and sequenced using GE108 and 377 primers (which are contained in Table 4). <sup>1</sup> [SEQ ID NO. 82-31] The mammalian expression vector N5KG4P is identical to the vector N5LG4P except it contains the human kappa light chain constant region in place of the human lambda light chain constant region. Sequencing of this 420 polynucleotide DNA revealed that it contains the entire kappa light chain variable domain.

A second independent PCR of the light chain variable region was performed using the 5' family 1 primer GE201 and the 3' primer GE204. (See primers for PCR of the <sup>1</sup> [SEQ ID NO. 5-2]) kappa light chain variable domain of 5E8 in Tables 1-3). The isolated PCR product was cloned into the TA vector (using methods previously described) and sequenced using Sp6 and T7 promoter primers. Sequencing revealed that this PCR product was identical to that obtained from the <sup>1</sup> [SEQ ID NO. 3] first PCR. The entire sequence of the light chain variable domain of primate monoclonal anti-human CD23 antibody 5E8 is presented below.

- 59 -

Light chain variable region of primate monoclonal antibody  
anti-human CD23 5E8

Leader

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu

ATG GAC ATG AGG GTC CCC GCT CAG CTC CTG GGG CTC CTT CTG CTC

-1

Trp Leu Pro Gly Ala Arg Cys

TGG CTC CCA GGT GCC AGA TGT

Mature Protein (Numbering is Kabat)

Framework 1

1

10

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val

GAC ATC CAG ATG ACC CAG TCT CCA TCT TCC CTG TCT GCA TCT GTA

20

23

Gly Asp Arg Val Thr Ile Thr Cys

GGG GAC AGA GTC ACC ATC ACT TGC

CDR 1

24

30

34

Arg Ala Ser Gln Asp Ile Arg Tyr Tyr Leu Asn

AGG GCA AGT CAG GAC ATT AGG TAT TAT TTA AAT

Framework 2

35

40

49

Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr

TGG TAT CAG CAG AAA CCA GGA AAA GCT CCT AAG CTC CTG ATC TAT

CDR2

50

56

Val Ala Ser Ser Leu Gln Ser

GTT GCA TCC AGT TTG CAA AGT

- 60 -

Framework 3

57                      60                      70  
Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Thr Glu Phe  
GGG GTC CCA TCA AGG TTC AGC GGC AGT GGA TCT GGG ACA GAG TTC

80

Thr Leu Thr Val Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr  
ACT CTC ACC GTC AGC AGC CTG CAG CCT GAA GAT TTT GCG ACT TAT

88

Tyr Cys  
TAC TGT

CDR 3

89    90                      97  
Leu Gln Val Tyr Ser Thr Pro Arg Thr  
CTA CAG GTT TAT AGT ACC CCT CGG ACG

Framework 4

98                      100                      107  
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
TTC GGC CAA GGG ACC AAG GTG GAA ATC AAA

2) Cloning the heavy chain variable domain of primate monoclonal anti-human CD23 antibody 5E8 by PCR

The first PCR of the heavy chain variable domain of 5E8 was performed using a set of 5' early leader heavy chain sequence primers and the 3' primer GE210. (See primers for PCR of the heavy chain variable domain of 6G5 and 5E8 in Table 1). A 420 base PCR product appeared in the family 3 primer reaction. The PCR product was puri-

[SEQ ID NOS: 5-9]  
^

B

fied and then digested with Nhe I and Sal I and cloned into the mammalian expression vector N5KG4P vector (as described previously). The PCR product was sequenced using the 268 and 928 primers. (See sequencing primers in Table 4.)  
[SEQ ID NOS.: 22-31]

A second independent PCR of the heavy chain variable domain of 5E8 was performed using the family 3 5' primer GE207 and the 3' primer GE210. (See primers for PCR of the heavy chain variable domain of 6G5 and 5E8 in Tables 1-3).  
[SEQ ID NOS.: 5-21]  
The isolated PCR product was cloned into a TA vector using the same techniques previously described and sequenced by using Sp6 and T7 primers. Sequencing revealed that the TAC at codon 91 had been changed into TGC.

In order to determine the appropriate codon at 91, a third independent PCR was performed using the same primers as the second PCR (see above). The PCR product was again cloned into a TA vector and sequenced using Sp6 and T7 primers. The sequence was found to be identical to the heavy chain variable sequence obtained in the first PCR. Therefore, the TGC at position 91 in the second independent PCR product is apparently the result of an error

introduced during PCR. This entire sequence of the heavy chain variable domain of primate monoclonal anti-human CD23 antibody <sup>5E8</sup> ~~6G5~~ is presented below.  
[SEQ ID NO.: 4]



- 62 -

Heavy chain variable region of primate monoclonal antibody  
anti-human CD23 SE8

Leader

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Pro Leu Leu Lys  
 ATG GAG TTT GGG CTG AGC TGG GTT TTC CTT GTT CCT CTT TTG AAA

-1

Gly Val Gln Cys  
 GGT GTC CAG TGT

Mature Protein (Numbering is Kabat)

Framework 1

1		10
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ala Lys Pro Gly		
GAG GTG CAG CTG GTG GAG TCT GGG GGC GGC TTG GCA AAG CCT GGG		
20		30
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Arg Phe Thr		
GGG TCC CTG AGA CTC TCC TGC GCA GCC TCC GGG TTC AGG TTC ACC		

CDR 1

31	35 35a 35b
Phe Asn Asn Tyr Tyr Met Asp	
TTC AAT AAC TAC TAC ATG GAC	

Framework 2

36	40	49
Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Val Ser		
TGG GTC CGC CAG GCT CCA GGG CAG GGG CTG GAG TGG GTC TCA		

```

50          52 52A  53                               60
Arg Ile Ser Ser Ser Gly Asp Pro Thr Trp Tyr Ala Asp Ser Val
CGT ATT AGT AGT AGT GGT GAT CCC ACA TGG TAC GCA GAC TCC GTG
65
Lys Gly
AAG GGC

```

```

56          70          80
Arg Phe Thr Ile Ser Arg Glu Asn Ala Asn Asn Thr Leu Phe Leu
AGA TTC ACC ATC TCC AGA GAG AAC GCC AAC AAC ACA CTG TTT CTT
      82 82a 82b 82c 83          90
Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
CAA ATG AAC AGC CTG AGA GCT GAG GAC ACG GCT GTC TAT TAC TGT
94
Ala Ser
GCG AGC

```

95 100 101  
Leu Thr Thr Gly Ser Asp Ser  
TTG ACT ACA GGG TCT GAC TCC

103						110		113
Trp	Gly	Gln	Gly	Val	Leu	Val	Thr	Ser
TGG	GGC	CAG	GGA	GTC	CTG	GTC	ACC	TCC

3) Construction of mammalian expression vectors

The heavy variable domain in N5KG4P was digested with Nhe I and Sal I, purified, and cloned into N5KG4P which contains the light chain variable domain of 5E8. This plasmid was then digested with the restriction endonucleases as previously described. This resulted in a vector containing both the light and heavy variable domain of 5E8. This vector was named N5KG4P + 5E8. The heavy and light variable domains of N5KG4P + 5E8 were then both inserted into the mammalian expression vector N5KG1 to create the N5KG1+ 5E8 vector.

4) Alteration of an amino acid in the heavy chain variable region of primate monoclonal antibody 5E8 by site specific mutagenesis and construction of mammalian expression vectors

Based upon the sequence of 5E8 heavy variable domain, there is a potential glycosylation site of the immunoglobulin at asparagine codon 75. This potential glycosylation site corresponds to a conserved asparagine-linked glycosylation motif having the following tripeptide sequence: (Asp) - (Any amino acid except proline) - (Serine or threonine). Therefore, a glycosylation mutant of 5E8, which would be unable to be glycosylated at this position because of modification of this glycosylation motif, was generated by replacing the asparagine codon 75 with a lysine (which is found in many human immunoglobulins at

- 65 -

this position). Site specific mutagenesis was effected by the following methods.

A first PCR was done using N5KG4P + 5E8 as a template and a 3' primer (corresponding to codon 71 to 79) and which contains a mutation at codon 75 (AAC changed to AAG, Primer MB1654, and a 5' primer at the beginning of the leader sequence (Primer MB1650). (See PCR Primers Used for the Generation of a Glycosylation Mutant of the Heavy

b Chain Variable Region 5E8 set forth in Table 5.) [SEQ ID NOS.: 32-35]

A second PCR was performed on the same template by using a 5' primer (corresponding to codon 71 to 79) containing the same mutation (Primer MB1653) and a 3' primer from the end of framework 4 (Primer MB1651) (See PCR Primers Used for the Generation of a Glycosylation Mutant

b of the Heavy Chain Variable Region of 5E8 in Table 5.) [SEQ ID NOS.: 32-35]

These two PCR products were isolated and mixed in equal molar ratios. A third independent PCR was then carried out by using the mixture of the first and second PCR products as a template with a 5' primer used in the first PCR (MB1650) and a 3' primer used in the second PCR (MB 1651). (See PCR Primers Used for the Generation of a Glycosylation Mutant of the Heavy Chain Variable Region in

b Table 5.) [SEQ ID NOS.: 32-35] The PCR product obtained in third PCR was found to contain the heavy variable domain coding region of 5E8 wherein the asparagine 75 had been changed to lysine.

The third PCR product was purified, digested with restriction endonucleases Sal I and Nhe I, and cloned into the N5KG4P containing the light variable domain only. The PCR product was sequenced to confirm that it comprised the mutant heavy variable domain. This mammalian expression plasmid was named N5KG4P + 5E8N-.

To construct of N5KG1 + 5E8N-, both light and heavy variable domains from N5KG4P - 5E8N- were digested with Bgl II and BsiW I, and Nhe I and Sal I respectively. The light variable domain was cloned first and then the heavy variable domain was inserted into the mammalian expression plasmid N5KG1, to create the plasmid N5KG1 + 5E8N-.

Table 1

Primers for PCR of the kappa light chain variable domain of 5E8

NAME	<u>Light chain Vκ -early leader 5' (Bgl II)</u>													FAMILY		
							-22	-21	-20	-19	-18	-17	-16	-15	-14	
GE201	5'	AT	CAC	<u>AGA TCT</u>	CTC	ACC	ATG	GAC	ATG	AGG	GTC	CCC	GCT	CAG	3'	1
GE200	5'	AT	CAC	<u>AGA TCT</u>	CTC	ACC			ATG	AGG	GTC	CCT	GCT	CAG	3'	2
GE202	5'	AT	CAC	<u>AGA TCT</u>	CTC	ACC			ATG	GAA	(A/G)CC	CAG	GC(T/G)	CAG		3'3
GE203	5'	AT	CAC	<u>AGA TCT</u>	CTC	ACC			ATG	GTG	TTG	CAG	ACC	CAG	GTC	3'4

Light chain Vk-3' primer (BsiW I)

					113	112	111	110	109	108	107	106	105	104	103
GE204	5'	GG	TGC	AGC	CAC	<u>CGT AGC</u>	TTT	GAT	(C/T)TC	CA(G/C)					
CTT	3'														

Table 2

Primers for PCR of the lambda light chain variable domain of 6G5

NAME	<u>Light chain VI-early leader 5' (Bgl II)</u>												FAMILY	
							-20	-19	-18	-17	-16	-15		
744	5'	AT	CAC	<u>AGA TCT</u>	CTC	ACC	ATG	(G/A)CC	TG(G/C)	TCC	CCT	CT	3'	1
745	5'	AT	CAC	<u>AGA TCT</u>	CTC	ACC	ATG	GCC	TGG	(A/G)CT	C(T/C)G	CT	3'	2
910	5'	AT	CAC	<u>AGA TCT</u>	CTC	ACC	ATG	GC(A/C)	TGG	A(T/C)C	CCT	CTC	3'	3

Light chain V1-3' primer (Avr II)

110 109 108 107 106 105 104  
926 5' (AC)10 CTT GGG CTG ACC TAG GAC GGT 3'

Table 3

Primers for PCR of the heavy chain  
variable domains from 6G5 and 5B8

NAME	Heavy chain-early leaders 5' (Sal I)	Family
	-20 -19 -18 -17 -16 -15	
MB1503	5' GCG ACT AAG TCG ACC ATG GAC TGG ACC TGG	3' 1
MB1502	5' GCG ACT AAG TCG ACC ATG AAA CAC CTG TGG	3' 2,4
GE207	5' GCG ACT AAG TCG ACC ATG GAG TTT GGG CTG AGC 3'	3
GE208	5' GCG ACT AAG TCG ACC ATG GGG TCA ACC GCC ATC 3'	5
GE209	5' GCG ACT AAG TCG ACC ATG TCT GTC TCC TTC CTC 3'	6

Heavy chain-3' primer (Nhe I)

120 119 118 117 116 115 114 113 112 111 110  
GE244 5' GC CAG GGG GAA GAC CGA TGG GCC CTT GGT GCT AGC TGA GGA GAC GG 3'  
GE210 5' GA TGG GCC CTT GGT GCT AGC TGA GGA GAC GG 3'

MB1533 5' GGT GCT AGC TGA GGA GAC GGT  
109 108 107 106 105 104 103 101 100 99  
GAC CAG GAC TCC CTG GCC CCA GAA GCC TAG 3'

Table 4

Sequencing Primers

Sp6 primer	5' AT TTA GGT GAC ACT ATA	3'
M13(-40)Forward Primer	5' GTT TTC CCA GTC ACG A	3'
T7 Promoter Primer	5' AT ATA CGA CTC ACT ATA GGG	3'
GE 108 Primer	5' CCG TCA GAT CGC CTG GAG ACG CCA 3'	
377 Primer	5' GCA GTT CCA GAT TTC AAC TG	3'
607 PRIMER	5' CCA GGC CAC TGT CAC GGC TTC	3'
266 PRIMER	5' CAG AGC TGG GTA CGT CCT CA	3'
268 PRIMER	5' GCC CCC AGA GGT GCT CTT GG	3'
876 PRIMER	5' ACA CAG ACC CGT CGA CAT GG	3'
928 PRIMER	5' GCT CTC GGA GGT GCT CCT GG	3'

Table 5

PCR Primers Used for the Generation of a Glycosylation Mutant of the Heavy Chain Variable Region of 5E8

		Sal I	-20	-19	-18	-17	-16	
MB 1650	5'	ACA GAC CCG	<u>TCG</u>	<u>ACC</u>	ATG GAG TTT GGG CTG	3'		
			Nhe I					
			118	117	116	115	114	113
MB 1651	5'	CCC CTT GGT	<u>GCT</u>	<u>AGC</u>	TGA GGA GAC GGT	3'		
			71	72	73	74	75	76
MB 1653	5'	AGA GAG AAC	GCC	AAG	AAC ACA CTG TTT	3'		
			79	78	77	76	75	74
MB 1654	5'	AAA CAG TGT GTT CTT	GGC	GTT	CTC TCT	3'		

C) Expression of PRIMATIZED® Antibodies in Chinese Hamster Ovary Cells

A large scale plasmid DNA was purified using the WIZARD® Maxipreps DNA Purification System (Promega Catalog # A7421).

The purified DNA was digested with Ssp I and BspLU11 I, precipitated with ethanol once, and resuspended in sterile TE.

Purified, endonuclease restricted plasmid DNA was then introduced into Chinese hamster ovary (CHO) dihydrofolate reductase minus DG44 cells using electroporation. The electroporation technique used is described below.

Approximately  $1.6 \times 10^6$  CHO cells were spun in an appropriate size sterile Corning tube for one minute at 1000 RPM. The media was removed and the cells were washed in fifteen milliliters of sterile ice cold SBS (sterile

sucrose buffered solution is 272 mM sucrose, 7 mM sodium phosphate pH 7.4, 1 mM  $MgCl_2$ ) and spun for 5 minutes at 1000 RPM. The SBS was removed and cells were suspended using fresh ice cold sterile SBS at a cell concentration of  $1 \times 10^7$  cells were per ml and left on ice for 15 minutes. The BTX 600 electroporator was turned on and preset at 230 volts, with the maximum voltage knobs being set at 500 volts/capacitance & resistance. The capacitance was set at 400 microfaradays and the resistance was set at 13 ohms (setting R1).

Plasmid DNA (4  $\mu g$  DNA or 2  $\mu g$  DNA) and 0.4 ml of cells ( $4 \times 10^6$  cells) were then placed in BTX 0.4 ml cuvettes (BTX Catalog # 620). The cells were shocked by placing the cuvette into the BTX 600 stand and pressing the automatic charge & pulse button. Approximately 20 separate electroporations were performed with each mammalian expression plasmid.

After shocking, the cuvettes were left at ambient temperature for fifteen minutes. The cells and DNA were from each cuvette were resuspended in 20 ml of CHO-SSFMII containing no hypoxanthine or thymidine (Gibco BRL Catalog # 31033-012) to which HT supplement (100X supplement is 10 mM sodium hypoxanthine, 1.6 mM thymidine Gibco BRL Catalog # 11067-014) had been added. The cells from a single electroporated cuvette were then plated into a 96 well plates (200  $\mu l$ /well) and placed into a 37°C CO<sub>2</sub> incubator.



Selection was started two or three days later by changing the media to the above media with the addition of 400 mg/ml of Geneticin® (G418, Gibco BRL Catalog # 10131-019). The cells were grown at 37°C and the cell media were changed every 3-5 days. After sixteen days G418 resistant clones appeared in the wells and the supernatant was assayed for antibody expression by ELISA. The highest expressing clones were then expanded individually. Monoclonal antibodies were purified as described below.

#### Immunoglobulin ELISA

Plates (Immulon 2, Dynatech Laboratories, Inc. Catalog # 011-010-3455) are coated overnight at 4°C with 200 ng unlabeled goat anti-human IgG antibody at 100 µl/well. This is effected using twenty milliliters of unlabeled goat anti-human IgG/10 mls Coating Buffer/plate (Boehringer Mannheim Ab Catalog # 605 400). (1:500 dilution of ~ 1mg/ml stock.) The coating buffer is then removed from the plates and dried using a paper towel. One hundred microliters of a dilution buffer/well is then added.

Antibody solutions and standards (100 ng/ml - 2.5 ng/ml) are then added in duplicate at 100 µl/well directly to the 100 ml dilution buffer. The antibody solutions and standards are contained in dilution buffer. The resultant solutions are then incubated for at least 1 hour at 37°C.

After incubation, the contents of each plate are removed and the plates are washed with tap water five times. The plates are then dried on a paper towel.

After drying of the plates, a second antibody is then added at 100  $\mu$ l/well. This second antibody is either goat anti-human Kappa-HRPO: added at 1/10,000 dilution or 1  $\mu$ l Ab/10 mls dilution buffer/plate, available from Southern Biotechnology Associates, Inc. Catalog # 2060-05 or a goat anti-human Lambda-HRPO; used at 1/20,000 dilution or 1  $\mu$ l Ab/20 mls dilution buffer/2 plates (available from Southern Biotechnology Associates, Inc. Catalog # 2070-05).

The antibody and contents of the plate are allowed to incubate for one hour at 37°C. After incubation the contents of each plate are removed. The plates are again washed five time with tap water, and the washed plates are dried. To the dried plates is then added HRPO substrate (TMB Microwell - two component) in an amount of 100  $\mu$ l/well. (Five milliliters of TMB Peroxidase Substrate + five milliliters of Peroxidase Solution B/plate (Kirdgaard and Perry Labs, TMB Microwell two component reagents Catalog # 50-76-00).

The reaction is stopped by the addition of one hundred microliters of 2M  $H_2SO_4$  to each well when the weakest standard (2.5 ng/ml) is visible over background. The optical density of wells in plates is then read using a plate reader, e.g., Molecular Devices Emax precision

- 72 -

microplate reader set at wavelength: OD 450 and OPT2 (OD 540).

### ELISA BUFFERS

#### Coating Buffer

Sodium Carbonate 0.8 gram / liter  
 Sodium Bicarbonate 1.55 gram / liter  
 Adjust pH to 9.5 with ~1ml 1N HCl

#### Dilution Buffer

0.5% Nonfat Dry Milk in PBS plus 0.01% Thimerosal  
 (5 gm/L) (100 mg/L)

Examples of ELISA values obtained using the above-described assay are set forth below.

<u>Standard</u>	<u>OD 450</u>	<u>OD450</u>	<u>Average</u>
100 ng/ml	0.805	0.876	0.841
50 ng/ml	0.395	0.472	0.434
25 ng/ml	0.213	0.252	0.233
10 ng/ml	0.089	0.105	0.097
5 ng/ml	0.054	0.055	0.055
2.5 ng/ml	0.031	0.035	0.033
0 ng/ml	0.004	0.006	0.005

### STANDARDS IN DILUTION BUFFER

Appropriate dilution of stock AB (sterile filtered in normal saline, protein determination by OD) to give 1mg/ml

**EXAMPLE:**

Chimeric monkey/human anti-CD4 (CE9.1) is 4.18 mg/ml  
24  $\mu$ l of above into 76  $\mu$ l Dilution Buffer is 1 mg/ml  
50  $\mu$ l Stock Ab (1 mg/ml) into 450  $\mu$ l Dilution Buffer  
(D.B.) is 100  $\mu$ g/ml  
50  $\mu$ l of above mixture into 450  $\mu$ l D.B. is 10  $\mu$ g/ml  
200  $\mu$ l of above mixture into 1.8 mls D.B. is 1  $\mu$ g/ml  
1 ml of above mixture into 9 mls D.B. is 100 ng/ml \*  
5 ml of above mixture into 5 ml D.B. is 50 ng/ml \*  
5 ml of above mixture into 5 ml D.B. is 25 ng/ml \*  
4 ml of above mixture into 6 ml D.B. is 10 ng/ml \*  
5 ml of above mixture into 5 ml D.B. is 5 ng/ml \*  
5 ml of above mixture into 5 ml D.B. is 2.5 ng/ml \*  
\* Standards used in the ELISA

**Antibody purification by Protein A****Procedure**

The culture supernatant is centrifuged to remove cells and debris. The centrifuge is then filtered through a 0.2 $\mu$ m filter. A protein A sepharose Fast flow column (recombinant protein A Sepharose Fast flow) (Pharmacia Biotech Catalog # 71-5000-09) is then prepared and equilibrated using PBS (pH 7.4).

The supernatant is loaded on the column at an appropriate flow rate (e.g., 2 ml/min). After loading, the column is washed with 10 column volume of PBS (pH 7.4).

The antibody is eluted from the column using an elution buffer (0.2M acetic acid, 0.1 M glycine pH 3.5) at 1 ml/min flow rate. One milliliter fractions/tube including 100  $\mu$ l of Tris are then collected. A spectrophotometer absorbance reading is then taken at 280 nm. The antibody fractions are then collected and dialyzed against PBS (pH 7.9) overnight. The dialysate is then sterilized by filtration through a 0.22  $\mu$ m membrane and stored at -20°C.

#### Assay Results

The PRIMATIZED® human gamma-4 anti-human CD23 antibodies which are described *supra*, were purified and assayed for induced IgE inhibitory activity *in vitro*. These results are contained in Figures 3 and 5. This was effected using the *in vitro* IL-4 IgE assay described *supra*.

These assay results surprisingly indicated that both human gamma-4 anti-human CD23 antibodies were not as active as the corresponding primate anti-human CD23 antibodies, i.e., they did not significantly inhibit induced IgE production *in vitro*.

However, because primate 5E8 and p5E8G4P have a potential asparagine linked glycosylation site in the heavy chain variable region, the effects of glycosylation at this site were investigated. (It was found that both these antibodies contain N-linked oligosaccharides at this site. (Data not shown.)) Therefore, in order to prevent

glycosylation, the asparagine in the glycosylation site was changed to a lysine in order to eliminate carbohydrate addition. This mutated antibody was named p5E8G4PN-. Assay results demonstrated that this antibody behaved identically to p5E8G4P in the IL-4 IgE assay (see Figure 3) and also exhibited an identical apparent affinity  $K_d$  for human CD23. (See Figure 4.) Therefore, these results indicated that the difference in IgE inhibition observed from the 5E8 gamma 4 PRIMATIZED® antibody in comparison to primate 5E8 antibody was not attributable to glycosylation differences.

The three primate antibodies (p5E8G4P, p5E8G4PN-, and pGG5G4P) were then expressed as human gamma-1 versions using substantially the same methodology. All three human gamma-1 anti-human CD23 antibodies, respectively designated p5E8G1, p5E8G1N- and pGG5G1, were found to be active in the *in vitro* IL-4/IgE assay (Figures 3 and 5).

p5E8G1 was found to be statistically more suppressive than p5E8G4P at a concentration of 0.3  $\mu\text{g/ml}$  ( $p[T,t]$  one tail + 0.0055) and at 3  $\mu\text{g/ml}$  ( $p[T<t]$  one tail + 0.0019). In addition, p5E8G1N- is statistically more suppressive than p5E8G4PN- at both 0.3  $\mu\text{g/ml}$  ( $p[T<t]$  one tail + 0.0392) and at 3  $\mu\text{g/ml}$  ( $p[T<t]$  one tail + 0.0310) (Figure 3).

- 76 -

Similarly, p6G5G1 completely inhibited induced IgE production at 3 mg/ml, while p6G5G4P did not. (These results are in Figure 5).

Thus, these results suggested that an active Fc region, in particular that of human gamma-1, is significant for induced IgE inhibition by anti-human CD23 antibodies.

#### EXAMPLE 2

To confirm our hypothesis as to the involvement of the Fc effector portion in IgE inhibition of anti-human CD23 antibodies, a third primate antibody, designated 2C8, also shown to inhibit IgE in in vitro) was converted to a F(ab')<sub>2</sub>. IgE inhibitory activity was determined using the same IL-4/IgE assay described previously.

#### Materials

The following materials were used in this example.

ImmunoPure F(ab')<sub>2</sub> Preparation Kit (Pierce Catalog # 44888)

digestion buffer: 20 mM sodium acetate buffer, pH 4.5

0.1 M citric acid, pH 3.0 (adjust pH with NaOH)

0.1% sodium azide in water

dialysis tubing; 50,000 MW cutoff (Spectra Por Catalog # 132 128)

shaking water bath capable of maintaining 37°C

- 77 -

polystyrene culture tubes, 17 x 100 mm (Fisher Catalog # 14-956-6B)

BCA protein assay (Pierce Catalog # 23224)

Centricon-50 concentrators (Amicon Catalog # 4225).

#### Equilibration of Immobilized Pepsin

0.25 milliliters of the 50% slurry of Immobilized Pepsin is added to a 17 x 100 mm test tube (0.125 ml of gel). Four milliliters of digestion buffer are then added. The pepsin is then separated from the buffer using the serum separator. The buffer is then discarded and the wash procedure repeated using another four milliliters of buffer. The immobilized pepsin is then resuspended in 0.5 ml of digestion buffer.

#### Preparation of Immobilized Protein A Column

Protein A AffinityPak® columns and ImmunoPure Binding and Elution buffers are brought to room temperature.

#### Preparation of 2C8 F(ab'), Fragments

2C8 F(ab')<sub>2</sub> fragments are prepared by methods well known in the antibody art. The inventors elected to use a commercially available kit, ImmunoPure F(ab')<sub>2</sub> Preparation Kit (Pierce Catalog # 44888), using the manufacturer's protocols.



Ten milligrams of lyophilized 2C8 antibody were dissolved in one milliliter of a digestion buffer (20 mM sodium acetate buffer, pH 4.5). One milliliter of the antibody containing sample was then added to a tube containing immobilized pepsin.

The antibody and immobilized pepsin were then incubated for four hours in a high speed shaker water bath at 37°C (at high speed), taking care to maintain the mixing constant during the incubation.

The resultant solubilized  $F(ab')_2$  and  $Fc$  fragments and the undigested IgGs were then recovered from the immobilized pepsin gel using a serum separator. The crude digest is then decanted into a clean tube.

In order to enhance recovery of  $F(ab')_2$  fragments, the immobilized pepsin desirably is then washed with 1.5 of milliliters of the ImmunoPure IgG binding buffer. The wash is then added to the crude digest.

The antibody fragments were then recovered using a protein A column. This is effected by opening an immobilized protein A column. Care is taken to avoid air bubbles from being drawn into the gel. The storage solution (which contains 0.02% sodium azide) is discarded.

The immobilized protein A column was then equilibrated using twelve milliliters of binding buffer (contained in ImmunoPure Preparation Kit). The column was then

transferred to a 17 x 100 mm test tube contained in the kit (labeled "F(ab')<sub>2</sub>") to collect eluate.

Three milliliters of the crude digest was then applied to a column and are allowed to flow completely into the gel. The use of AffinityPak™ columns is desirable as these columns stop flowing automatically when the level reaches the top frit.

The column is then washed using six milliliters of binding buffer. The eluate which contains F(ab')<sub>2</sub> fragments was then collected. This eluate also contains small Fc fragments that can no longer bind protein A (which are not bound to the Protein A column). However, the substantial portion thereof was eliminated by dialysis.

Dialysis was effected by taking the F(ab')<sub>2</sub> containing eluate and dialyzing the eluate against pH 7.4 phosphate buffered saline, using dialysis tubing with a molecular weight cut-off of 50,000 so as to eliminate the small Fc fragment containments (Spectra Pur. Catalog # 132 128).

This resulted in a F(ab')<sub>2</sub> fraction having an optical density of 280 nm of 0.707 (6 ml). After dialysis and concentration with Centricon-50 concentrators (Amicon Catalog # 4225), the 2C8 F(ab')<sub>2</sub> product was assayed for protein content using a BCA protein assay (Pierce Catalog # 23224). The protein content was found to be 3.76 mg per milliliter.

- 80 -

The 2C8 F(ab)<sub>2</sub>'s were assayed for IgE inhibitory activity and were found to be substantially incapable of inhibiting IgE production in the same *in vitro* assays described previously. These results are contained in Figure 6. In fact, the F(ab')<sub>2</sub> was found to antagonize the suppressive effects of induced IgE on the monoclonal antibody 2C8. These results are in Figure 7.

### EXAMPLE 3

The PRIMATIZED® gamma 1 and gamma 4P versions of primate monoclonal 6G5 were both evaluated for their effect on inhibition of induced IgE production *in vivo* in the SCID mouse model described previously. p5E8G1N- was found to be as efficient as primate 5E8 in inhibiting induced IgE. (See Figures 8 and 9). While neither provide 6G5 nor the primatized p6G5G4P were effective at inhibiting induced IgE *in vivo*, primatized p6G5G1 inhibited induced IgE production. (See Figures 9 and 10.) These results further substantiate our conclusion that an active Fc region is significant to the ability of an anti-human CD23 antibody to effectively inhibit induced IgE production.

### Utility

The subject anti-human CD23 antibodies which comprise human gamma-1 constant domains, because of their ability

to effectively inhibit IgE production, are effective in treating any disease wherein inhibition of IgE production is therapeutically desirable. Such diseases include by way of example allergic diseases, autoimmune diseases and inflammatory disease.

Specific conditions which are potentially treatable by administration of the subject anti-CD23 human gamma-1 constant domain containing antibodies include the following:

Allergic bronchopulmonary aspergillosis; Allergic rhinitis and conjunctivitis autoimmune hemolytic anemia; Acanthosis nigricans; Allergic contact dermatitis; Addison's disease; Atopic dermatitis; Alopecia areata; Alopecia universalis; Amyloidosis; Anaphylactoid purpura; Anaphylactoid reaction; Aplastic anemia; Angioedema, hereditary; Angioedema, idiopathic; Ankylosing spondylitis; Arteritis, cranial; Arteritis, giant cell; Arteritis, Takayasu's; Arteritis, temporal; Asthma; Ataxia-telangiectasia; Autoimmune oophoritis; Autoimmune orchitis; Autoimmune polyendocrine failure; Behcet's disease; Berger's disease; Buerger's disease; bronchitis; Bullous pemphigus; Candidiasis, chronic mucocutaneous; Caplan's syndrome; Post-myocardial infarction syndrome; Post-pericardiotomy syndrome; Carditis; Celiac sprue; Chagas's disease; Chediak-Higashi syndrome; Churg-Strauss disease; Cogan's syndrome; Cold agglutinin disease; CREST

syndrome; Crohn's disease; Cryoglobulinemia; Cryptogenic fibrosing alveolitis; Dermatitis herpetiformis; Dermatomyositis; Diabetes mellitus; Diamond-Blackfan syndrome; DiGeorge syndrome; Discoid lupus erythematosus; Eosinophilic fasciitis; Episcleritis; Drythema elevatum diutinum; Erythema marginatum; Erythema multiforme; Erythema nodosum; Familial Mediterranean fever; Felty's syndrome; Fibrosis pulmonary; Glomerulonephritis, anaphylactoid; Glomerulonephritis, autoimmune; Glomerulonephritis, post-streptococcal; Glomerulonephritis, post-transplantation; Glomerulopathy, membranous; Goodpasture's syndrome; Graft-vs.-host disease; Granulocytopenia, immune-mediated; Granuloma annulare; Granulomatosis, allergic; Granulomatous myositis; Grave's disease; Hashimoto's thyroiditis; Hemolytic disease of the newborn; Hemochromatosis, idiopathic; Henoch-Schoenlein purpura; Hepatitis, chronic active and chronic progressive; Histiocytosis\_X; Hypereosinophilic syndrome; Idiopathic thrombocytopenic purpura; Job's syndrome; Juvenile dermatomyositis; Juvenile rheumatoid arthritis (Juvenile chronic arthritis); Kawasaki's disease; Keratitis; Keratoconjunctivitis sicca; Landry-Guillain-Barre-Strohl syndrome; Leprosy, lepromatous; Loeffler's syndrome; lupus; Lyell's syndrome; Lyme disease; Lymphomatoid granulomatosis; Mastocytosis, systemic; Mixed connective tissue disease; Mononeuritis multiplex; Muckle-Wells syndrome; Mucocutane-

ous lymph node syndrome; Mucocutaneous lymph node syndrome; Multicentric reticulohistiocytosis; Multiple sclerosis; Myasthenia gravis; Mycosis fungoides; Necrotizing vasculitis, systemic; Nephrotic syndrome; Overlap syndrome; Panniculitis; Paroxysmal cold hemoglobinuria; Paroxysmal nocturnal hemoglobinuria; Pemphigoid; Pemphigus; Pemphigus erythematosus; Pemphigus foliaceus; Pemphigus vulgaris; Pigeon breeder's disease; Pneumonitis, hypersensitivity; Polyarteritis nodosa; Polymyalgia rheumatic; Polymyositis; Polyneuritis, idiopathic; Portuguese familial polyneuropathies; Pre-eclampsia/eclampsia; Primary biliary cirrhosis; Progressive systemic sclerosis (Scleroderma); Psoriasis; Psoriatic arthritis; Pulmonary alveolar proteinosis; Pulmonary fibrosis, Raynaud's phenomenon/syndrome; Reidel's thyroiditis; Reiter's syndrome, Relapsing polychondritis; Rheumatic fever; Rheumatoid arthritis; Sarcoidosis; Scleritis; Sclerosing cholangitis; Serum sickness; Sezary syndrome; Sjogren's syndrome; Stevens-Johnson syndrome; Still's disease; Subacute sclerosing panencephalitis; Sympathetic ophthalmia; Systemic lupus erythematosus; Transplant rejection; Ulcerative colitis; Undifferentiated connective tissue disease; Urticaria, chronic; Urticaria, cold; Uveitis; Vitiligo; Weber-Christian disease; Wegener's granulomatosis; Wiskott-Aldrich syndrome.

Of these, the preferred indications treatable or presentable by administration of anti-CD23 antibodies include allergic rhinitis and conjunctivitis, atopic dermatitis; eczema; Job's syndrome, asthma; and allergic conditions; chronic inflammatory diseases and conditions.

The amount of antibody useful to produce a therapeutic effect can be determined by standard techniques well known to those of ordinary skill in the art. The antibodies will generally be provided by standard technique within a pharmaceutically acceptable buffer, and may be administered by any desired route. Because of the efficacy of the presently claimed antibodies and their tolerance by humans it is possible to administer these antibodies repetitively in order to combat various diseases or disease states within a human.

One skilled in the art would be able, by routine experimentation, to determine what an effective, non-toxic amount of antibody would be for the purpose of effecting allergic diseases and inflammatory conditions. Generally, however, an effective dosage will be in the range of about 0.05 to 100 milligrams per kilogram body weight per day.

The antibodies of the invention may be administered to a human or other animal in accordance with the aforementioned methods of treatment in an amount sufficient to produce such effect to a therapeutic or prophylactic degree. Such antibodies of the invention can be adminis-

tered to such human or other animal in a conventional dosage form prepared by combining the antibody of the invention with a conventional pharmaceutically acceptable carrier or diluent according to known techniques. It will be recognized by one of skill in the art that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables.

The route of administration of the antibody of the invention may be oral, parenteral, by inhalation or topical. The term parenteral as used herein includes intravenous, intramuscular, subcutaneous, rectal, vaginal or intraperitoneal administration. The intravenous form of parenteral administration is generally preferred.

The daily parenteral and oral dosage regimens for employing compounds of the invention to prophylactically or therapeutically induce immunosuppression will generally be in the range of about 0.05 to 100, but preferably about 0.5 to 10, milligrams per kilogram body weight per day.

The antibody of the invention may also be administered by inhalation. By "inhalation" is meant intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques. The preferred dosage amount of a



compound of the invention to be employed is generally within the range of about 10 to 100 milligrams.

The antibody of the invention may also be administered topically. By topical administration is meant non-systemic administration and includes the application of an antibody (or fragment thereof) compound of the invention externally to the epidermis, to the buccal cavity and instillation of such an antibody into the ear, eye and nose, and where it does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal and intramuscular administration. The amount of an antibody required for therapeutic or prophylactic effect will, of course, vary with the antibody chosen, the nature and severity of the condition being treated and the animal undergoing treatment, and is ultimately at the discretion of the physician. A suitable topical dose of an antibody of the invention will generally be within the range of about 1 to 100 milligrams per kilogram body weight daily.

#### Formulations

While it is possible for an antibody or fragment thereof to be administered alone, it is preferable to present it as a pharmaceutical formulation. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, e.g., from 1% to 2% by weight of the

formulation, although it may comprise as much as 10% w/w but preferably not in excess of 5% w/w and more preferably from 0.1% to 1% w/w of the formulation.

The topical formulations of the present invention, comprise an active ingredient together with one or more acceptable carrier(s) therefor and optionally any other therapeutic ingredients(s). The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of where treatment is required, such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 90°-100°C for half an hour. Alternatively, the solution may be steril-

ized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol and propylene glycol.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or

olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or macrogels. The formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surface active such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

It will be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of an antibody or fragment thereof of the invention will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular animal being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of an antibody or fragment thereof of the invention given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The

following are, therefore, to be construed as merely illustrative examples and not a limitation of the scope of the present invention in any way.

#### Capsule Composition

A pharmaceutical composition of this invention in the form of a capsule is prepared by filling a standard two-piece hard gelatin capsule with 50 mg. of an antibody or fragment thereof of the invention, in powdered form, 100 mg. of lactose, 32 mg. of talc and 8 mg. of magnesium stearate.

#### Injectable Parenteral Composition

A pharmaceutical composition of this invention in a form suitable for administration by injection is prepared by stirring 1.5k by weight of an antibody or fragment thereof of the invention in 10k by volume propylene glycol and water. The solution is sterilized by filtration.

#### Ointment Composition

Antibody or fragment thereof of the invention 10 g.  
White soft paraffin to 100.0 g.

The antibody or fragment thereof of the invention is dispersed in a small volume of the vehicle to produce a smooth, homogeneous product. Collapsible metal tubes are then filled with the dispersion.

Topical Cream Composition

Antibody or fragment thereof of the invention 1.0g.

Polawax GP 200 20.0 g.

Lanolin Anhydrous 2.0 g.

White Beeswax 2.5 g.

Methyl hydroxybenzoate 0.1 g.

Distilled Water to 100.0 g.

The polawax, beeswax and lanolin are heated together at 60°C. A solution of methyl hydroxybenzoate is added and homogenization is achieved using high speed stirring. The temperature is then allowed to fall to 50°C. The antibody or fragment thereof of the invention is then added and dispersed throughout, and the composition is allowed to cool with slow speed stirring.

Topical Lotion Composition

Antibody or fragment thereof of the invention 1.0g.

Sorbitan Monolaurate 0.6 g. Polysorbate 20 0.6 g.

Cetostearyl Alcohol 1.2 g. Glycerin 6.0 g.

Methyl Hydroxybenzoate 0.2 g.

Purified Water B.P. to 100.00 ml. (B.P. = British Pharmacopeia)

The methyl hydroxybenzoate and glycerin are dissolved in 70 ml. of the water at 75°C. The sorbitan monolaurate, polysorbate 20 and cetostearyl alcohol are melted together at 75°C and added to the aqueous solution. The resulting

- 92 -

emulsion is homogenized, allowed to cool with continuous stirring and the antibody or fragment thereof of the invention is added as a suspension in the remaining water. The whole suspension is stirred until homogenized.

**Eye Drop Composition**

Antibody or fragment thereof of the invention 0.5g.

Methyl Hydroxybenzoate 0.01 g.

Propyl Hydroxybenzoate 0.04 g.

Purified Water B.P. to 100.00 ml.

The methyl and propyl hydroxybenzoates are dissolved in 70 ml. purified water at 75°C and the resulting solution is allowed to cool. The antibody or fragment thereof of the invention is then added, and the solution is sterilized by filtration through a membrane filter (0.022  $\mu$ m pore size), and packed aseptically into suitable sterile containers.

**Composition for Administration by Inhalation**

For an aerosol container with a capacity of 15-20 ml: mix 10 mg. of an antibody or fragment thereof of the invention with 0.2-0.5k of a lubricating agent, such as polysorbate 85 or oleic acid, and disperse such mixture in a propellant, such as freon, preferably in a combination of (1,2 dichlorotetrafluoroethane) and difluorochloromethane and put into an appropriate aerosol container adapted

- 93 -

for either intranasal or oral inhalation administration. Composition for Administration by Inhalation For an aerosol container with a capacity of 15-20 ml: dissolve 10 mg. of an antibody or fragment thereof of the invention in ethanol (6-8 ml.), add 0.1-0.2k of a lubricating agent, such as polysorbate 85 or oleic acid; and disperse such in a propellant, such as freon, preferably in combination of (1-2 dichlorotetrafluoroethane) and difluorochloromethane, and put into an appropriate aerosol container adapted for either intranasal or oral inhalation administration.

**Parenteral Administrable Antibody Compositions**

The antibodies and pharmaceutical compositions of the invention are particularly useful for parenteral administration, i.e., subcutaneously, intramuscularly or intravenously. The compositions for parenteral administration will commonly comprise a solution of an antibody of the invention or a cocktail thereof dissolved in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be employed, e.g., water, buffered water, 0.4k saline (normal saline), 0.3% glycine, and the like. The use of normal saline is preferred. These solutions are sterile and generally free of particulate matter. These solutions may be sterilized by conventional, well-known sterilization techniques. The compositions may contain pharmaceutically acceptable auxiliary sub-



stances as required to approximate physiological conditions such as pH adjusting and buffering agents, etc. The concentration of the antibody or fragment thereof of the invention in such pharmaceutical formulation can vary widely. Such concentrations will be selected primarily based on fluid volumes, viscosities, etc., according to the particular mode of administration selected. Generally suitable intravenous concentrations range from about one to one hundred milligrams per milliliter.

Thus, a pharmaceutical composition of the invention for intravenous injection could comprise 10 mL normal saline containing 40-50 mg of an anti-human CD23 antibody of the invention. Methods for preparing parenterally administrable compositions are well-known or will be apparent to those skilled in the art, and are described in more detail in, for example, *Remington's Pharmaceutical Science*, 15th ed., Mack Publishing Company, Easton, Pennsylvania, hereby incorporated by reference herein.

The antibodies of the invention can be lyophilized for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with conventional immune globulins and art-known lyophilization and reconstitution techniques can be employed.

Depending on the intended result, the pharmaceutical composition of the invention can be administered for prophylactic and/or therapeutic treatments. In therapeu-

tic application, compositions are administered to a patient already suffering from a disease, in an amount sufficient to cure or at least partially arrest the disease and its complications. In prophylactic applications, compositions containing the present antibodies or a cocktail thereof are administered to a patient not already in a disease state to enhance the patient's resistance.

Single or multiple administrations of the pharmaceutical compositions can be carried out with dose levels and pattern being selected by the treating physician. In any event, the pharmaceutical composition of the invention should provide a quantity of the subject anti-CD23 antibodies sufficient to effectively treat the patient.

It should also be noted that the antibodies of this invention may be used for the design and synthesis of either peptide or non-peptide compounds (mimetics) which would be useful in the same therapy as the antibody. See, e.g., Saragovi et al., *Science*, 253:792-795 (1991).

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without diverting from the scope of the invention. Accordingly, the invention is not limited by the appended claims.